

Objectives

Induction and Recursion

Dr. Mattox Beckman

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN
DEPARTMENT OF COMPUTER SCIENCE

- ▶ Identify the parts of a proof by induction and their corresponding parts in a recursive function.
- ▶ Identify the requirements for a recursive function to terminate with a correct answer.

Induction

A proof by induction works by making two steps do the work of an infinite number of steps. It's really a way of being very lazy!

- ▶ Pick a property $P(n)$ which you'd like to prove for all n .
- ▶ **Base case:** Prove $P(n)$, for $n = 1$, or whatever n 's smallest value should be.
- ▶ **Induction case:** You want to prove $P(n)$, for all n . To do that, *assume* that $P(n - 1)$ is true, and use that information to prove that $P(n)$ has to be true.

The idea is that there are an infinite number of n such that $P(n)$ is true. But with this technique you only had to prove two cases.

Induction Example

To prove: Let $P(n) =$ "The sum of the first n odd numbers is n^2 ."

Induction Example

To prove: Let $P(n)$ = "The sum of the first n odd numbers is n^2 ."
Base case: Let $n = 1$. Then $n^2 = 1$, and the sum of the list $\{1\}$ is 1; therefore the base case holds.

Induction Example

To prove: Let $P(n)$ = "The sum of the first n odd numbers is n^2 ."
Base case: Let $n = 1$. Then $n^2 = 1$, and the sum of the list $\{1\}$ is 1; therefore the base case holds.
Induction case: Suppose you need to show that this property is true for some n . First, pretend that somebody else already did all the work of proving that $P(n - 1)$ is true. Now use that to show that $P(n)$ is true, and take all the credit.

Induction Example

To prove: Let $P(n)$ = "The sum of the first n odd numbers is n^2 ."
Base case: Let $n = 1$. Then $n^2 = 1$, and the sum of the list $\{1\}$ is 1; therefore the base case holds.
Induction case: Suppose you need to show that this property is true for some n . First, pretend that somebody else already did all the work of proving that $P(n - 1)$ is true. Now use that to show that $P(n)$ is true, and take all the credit.

$$(1 + 3 + 5 + \dots + 2n - 3) = (n - 1)^2$$

So add $2n - 1$ to both sides ...

$$\Rightarrow (1 + 3 + 5 + \dots + 2n - 3 + 2n - 1) = (n - 1)^2 + 2n - 1$$
$$\Rightarrow n^2 - 2n + 1 + 2n - 1$$
$$\Rightarrow n^2$$

Recursion

A recursive routine has a similar structure. You have a base case, a recursive case, and a conditional to check which case is appropriate.

- ▶ Pick a function $f(n)$ which you'd like to compute for all n .
- ▶ **Base case:** Compute $f(n)$, for $n = 1$, or whatever n 's smallest value should be.
- ▶ **Recursive case:** Assume that someone else already computed $f(n - 1)$ for you. Use that information to compute $f(n)$, and then take all the credit.

Iterating Recursion Example

Suppose you want a recursive routine that computes the *n*th square.

```
nthsq 1 = 1
nthsq n = 2*n-1 + nthsq (n-1)
```

- ▶ The pattern matching checks which case is appropriate.
- ▶ Line 1 is the base case – it stops the recursion.
- ▶ Line 2 is the recursive case.

Important Things about Recursion

```
nthsq 1 = 1
nthsq n = 2*n-1 + nthsq (n-1)
```

- ▶ Your base case has to stop the computation.
- ▶ Your recursive case has to call the function with a *smaller* argument than the original call.
- ▶ Your *if* statement has to be able to tell when the base case is reached.
- ▶ Failure to do any of the above will cause an infinite loop.

History

(Discovered on Wikipedia)

- ▶ The proof that that the first *n* odd numbers sums to *n*² first appeared in *Arithmeticonum libri duo* by Francesco Maurolico in 1575.
- ▶ Wikipedia says it's the earliest known *explicit* use of proof by induction.
- ▶ Implicit uses of proof by induction can be found in the writings of Plato and Euclid in the 300's BCE.

